

Electrolysis of potassium iodide solution

Filter paper soaked in potassium iodide solution which also contains starch and phenolphthalein is placed on an aluminium sheet which forms one electrode of an electric circuit. The other electrode is used as a 'pen nib' to 'write' on the filter paper. When this electrode is made positive and the aluminium sheet negative, the writing is blue/black and when the polarity is reversed, the writing is pink.

This experiment should take around 20 minutes.

Apparatus and chemicals

- Eye protection
- Disposable gloves
- Aluminium (or other metal) sheet approximately 25 cm × 25 cm but the size is not critical
- DC power supply (0 - 12 V)
- Leads and crocodile clips to connect to the power pack
- Potassium iodide solution (**Low hazard**) (0.25 mol dm^{-3} , but the concentration is not critical) (100 cm^3)
- Starch solution (20 cm^3) (see *Technical notes* below for how to prepare)
- Sodium thiosulfate solution (**Low hazard**) (approximately 1 mol dm^{-3} , but the concentration is not critical) (a few drops)
- Phenolphthalein solution (20 cm^3) (**Highly flammable**) (see *Technical notes* below for how to prepare)
- Filter papers, as large as possible to fit on the aluminium sheet

Technical notes

Aqueous potassium iodide (**Low hazard**) Refer to SSERC or CLEAPSS Hazcard Sodium thiosulfate solution (**Low hazard**)

Phenolphthalein solution (**Highly flammable**) Refer to SSERC or CLEAPSS Recipe and Hazcards

Procedure

HEALTH & SAFETY: Both demonstrator and audience should wear eye protection.

Wash hands after the experiment. Demonstrators with skin problems or cuts should wear disposable gloves.

a Mix 40 cm^3 of potassium iodide solution with 10 cm^3 of starch solution. If the resulting solution has a blue colour (caused by contamination with iodine) add sodium thiosulfate solution dropwise until the solution becomes colourless. Then add 10 cm^3 of phenolphthalein solution. If the resulting solution has a pink colour (caused by contamination with alkali) add dilute hydrochloric acid solution dropwise until the solution just becomes colourless.

b Thoroughly moisten three sheets of filter paper in this prepared mixture and place the papers onto the aluminium sheet one on top of the other. Moistening may be done using a dropping pipette or wash bottle. Connect the aluminium sheet to the negative terminal of the power supply using a lead and crocodile clip. Connect a second lead to the positive terminal and switch on the power pack at



between 6 V and 12 V. Now use the end of the positive lead to write or draw something on the top sheet of filter paper. The writing will appear blue/black, as iodine is discharged at the positive electrode and reacts with the starch to produce a blue/black complex. A corresponding pink line will appear on the lower filter paper in contact with the aluminium sheet but this will not be visible. This is caused by the discharge of H^+ ions as hydrogen which leaves an excess of OH^- ions in the solution. This alkaline solution turns the phenolphthalein pink.

c Then switch the polarity of the electrodes so that the aluminium sheet becomes positive, and the free lead negative. The writing on the upper sheet of filter paper now becomes pink. (There will also be a corresponding blue/black line on the lower filter paper.) The reason for using three filter papers in a stack is that this blue/black line would be visible through a single moist filter paper and obscure the paler pink line. Some teachers may wish to draw attention to the lines underneath the filter papers, other may wish to ignore them.

Extensions

If alternating current is used, a dotted line of alternating pink and blue/black is seen provided the lead is drawn over the filter paper quickly enough. This is not as spectacular as might be expected as the pink is much paler than the blue/black.

Indicators other than phenolphthalein could be tried.

Theory

An aqueous solution of potassium iodide contains the following ions:

$\text{K}^+(\text{aq})$ and $\text{I}^-(\text{aq})$ from the solute $\text{H}^+(\text{aq})$ and $\text{OH}^-(\text{aq})$ from dissociation of the water.

At the positive electrode

$\text{I}^-(\text{aq})$ and $\text{OH}^-(\text{aq})$ are attracted to the positive electrode (anode) where iodide ions are converted to iodine: $2\text{I}^-(\text{aq}) - 2\text{e}^- \rightarrow \text{I}_2(\text{aq})$

(This occurs in preference to the thermodynamically favoured

$4\text{OH}^-(\text{aq}) - 4\text{e}^- \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$ because of the much higher concentration of $\text{I}^-(\text{aq})$ than $\text{OH}^-(\text{aq})$).

This iodine then forms a blue/black complex with the starch.

At the negative electrode

$\text{H}^+(\text{aq})$ and $\text{K}^+(\text{aq})$ are attracted to the negative electrode (cathode) where hydrogen ions are converted to hydrogen: $2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$

(This occurs in preference to $\text{K}^+(\text{aq}) + \text{e}^- \rightarrow \text{K}(\text{s})$ because the discharge potential is more positive.)

This leaves an excess of $\text{OH}^-(\text{aq})$ ions around the cathode which turn phenolphthalein pink.

Teaching notes

A simpler explanation as to why hydrogen is discharged at the cathode is that if potassium (the alternative product) were discharged it would immediately react with water to return to K^+ ions, OH^- ions and hydrogen. Teachers might wish to use this explanation with some students.

A sheet of newspaper placed below the aluminium sheet will reduce mess if the filter papers have been over-enthusiastically moistened.



This experiment is suitable for a class experiment or science club activity if sufficient apparatus is available - a biscuit tin lid (or almost any metal sheet) can be used as an alternative to the aluminium sheet.

Attaching a graphite pencil sharpened at both ends to the second lead gives a good, narrow line and also nicely shows the conductivity of graphite at the same time.

Technical notes

The starch solution must be prepared shortly before use – it will not keep. It is prepared by mixing 1 g of soluble starch with a little deionised water to form a thin paste then adding to this paste 80 cm³ of boiling water. Stir the mixture, allow it to cool and dilute to 100 cm³.

To make phenolphthalein solution, dissolve 1 g solid phenolphthalein in 600 cm³ of industrial methylated spirits and make up to 1 dm³ with water.

Reference

This experiment has been adapted by the RSC

The RSC wishes to thank Dr Vladimir Volkovich formerly of Manchester University for translating an early draft of this demonstration from Russian. It also thanks Vanessa Byrne and Alison Oliver of North Leamington School for providing laboratory facilities and technical assistance. Dr Lynn Nickerson of Didcot Girls' School trialled the experiment her Science Club, which made a number of helpful suggestions.

Credits

© Royal Society of Chemistry

Health & safety checked January 2018

Page last updated March 2018

