

# Scaffolding explanations

## Target level

These materials are designed for students taking post-16 chemistry courses.

## Topics

The topics covered in these materials include charge: size ratio (charge density); atomic structure and core charge; electronegativity; bond polarity; lattice energy; hydration of ions; hydrogen bonding; melting temperature; boiling temperature; atomic size and base strength.

## Rationale

These materials are designed to support students in developing confidence in using explanations in chemistry. In particular the materials aim to demonstrate how a range of phenomena that chemists study are explained in terms of a limited set of basic chemical concepts. The materials are designed to support students by providing frameworks within which they can successfully complete explanations, as a step towards developing explanations on their own. The materials also require students to translate information between a schematic form and standard prose. These notions of 'scaffolding' students' learning, and providing active learning tasks, are discussed in Chapters 3 and 5 of the Teachers' notes.

The materials comprise of two sets of questions (of a type commonly used in public examinations) requiring explanations - which may be used as a pre-test and post-test - a worksheet explaining the key concepts related to these questions, and a worksheet providing structured support in working through the first set of questions.

During piloting it was found some students found the materials helpful, and it was suggested that similar exercises in a wider range of topics would be useful. Some students found the questions difficult, but others found the materials repetitive. This latter response could be seen as a positive outcome, as clearly the students were realising that a few key ideas could be used in a wide range of situations. However, teachers may also wish to use the pre-test to distinguish those students who are already able to use the key concepts to develop satisfactory explanations from those who would benefit from working through the complete set of materials.

## Instructions

Each student requires a copy of the following worksheets

- **Explaining chemical phenomena (1)** – which may be used as a pre-test
- **Constructing chemical explanations** – which provides a review of the key ideas
- **Completing explanations** – which provides support in answering the questions on the pre-test
- **Explaining chemical phenomena (2)** – which may be used as a post-test

## Resources

- Student worksheets
  - Explaining chemical phenomena (1)
  - Constructing chemical explanations

## RS•C

- Completing explanations
- Explaining chemical phenomena (2)
- Examples of chemical explanations (1) and (2) – (answers to Explaining chemical phenomena (1) and (2))

### Feedback for students

A teachers' answer sheet for **Constructing chemical explanations** is provided. The completed schematics for the questions in the pre- and post- tests are also provided as **Examples of chemical explanations** (1) and (2).

# Scaffolding explanations – answers

## Constructing chemical explanations

The table is reproduced below. The answers have been printed in **bold**.

E	Z	$n_p$	$n_e$	e.c.	$n_{ve}$	$n_{ce} (=n_e - n_{ve})$	$n_p - n_{ce} =$	c.c.
H	1	1	1	1	1	0	1-0 =	+1
He	2	2	2	2	2	0	2-0 =	+2
Li	3	3	3	2.1	1	2	3-2 =	+1
Be	4	4	4	2.2	2	2	4-2 =	+2
B	5	5	5	2.3	3	2	5-2 =	+3
C	6	6	6	2.4	<b>4</b>	2	<b>6-2 =</b>	+4
N	7	7	7	<b>2.5</b>	5	2	7-2 =	+5
O	8	8	8	2.6	6	2	8-2 =	<b>+6</b>
F	9	9	9	2.7	<b>7</b>	<b>2</b>	<b>9-2 =</b>	+7
Ne	10	10	10	2.8	<b>8</b>	<b>2</b>	10-2 =	<b>+8</b>
Na	11	11	11	2.8.1	1	10	11-10 =	+1
Mg	<b>12</b>	<b>12</b>	<b>12</b>	2.8.2	2	10	12-10 =	+2
Al	13	13	13	2.8.3	3	10	<b>13-10 =</b>	+3
Si	14	<b>14</b>	<b>14</b>	2.8.4	4	<b>10</b>	<b>14-10 =</b>	+4
P	<b>15</b>	<b>15</b>	<b>15</b>	2.8.5	5	10	<b>15-10 =</b>	<b>+5</b>
S	<b>16</b>	<b>16</b>	<b>16</b>	<b>2.8.6</b>	6	10	<b>16-10 =</b>	<b>+6</b>
Cl	<b>17</b>	<b>17</b>	<b>17</b>	2.8.7	7	<b>10</b>	<b>17-10 =</b>	+7
Ar	18	18	18	2.8.8	8	10	18-10 =	+8
K	19	<b>19</b>	<b>19</b>	2.8.8.1	1	18	19-18 =	<b>+1</b>
Ca	20	20	20	<b>2.8.8.2</b>	2	<b>18</b>	<b>20-18 =</b>	+2

Completing the core charge diagram

B: +3    O: +6

P: +5    S: +6

K: +1


(the core charge matches the element's position across the period)

Other answers are provided as photocopyable masters.


# Examples of chemical explanations (1)

The following scheme may be used as the basis of explaining the eight questions included in the worksheets **Explaining chemical phenomena** (1).


1. Lithium has a higher melting temperature (454 K) than sodium (371 K).




2. There is stronger bonding, called hydrogen bonding, between molecules of water ( $\text{H}_2\text{O}$ ) than between molecules of hydrogen sulfide ( $\text{H}_2\text{S}$ ).




3. The nitrogen atom is smaller than the carbon atom (ie it has a smaller covalent radius - 0.074 nm compared to 0.077 nm).



4. Chlorine is more electronegative than bromine.




5. Ammonia ( $\text{NH}_3$ ) is a stronger base than phosphine ( $\text{PH}_3$ ).







6. Magnesium chloride has a larger lattice energy ( $2489 \text{ kJ mol}^{-1}$ ) than calcium chloride ( $2197 \text{ kJ mol}^{-1}$ ).



7. More energy is released when sodium ions are hydrated ( $390 \text{ kJmol}^{-1}$ ) than when potassium ions are hydrated ( $305 \text{ kJmol}^{-1}$ ).




8. Nitrogen is less electronegative than oxygen.




# Examples of chemical explanations (2)

The following schemes may be useful as the basis for explaining the eight included in the worksheet **Explaining chemical phenomena (2)**


1. Potassium has a lower melting temperature (336 K) than sodium (371 K).




2. There is weaker bonding between molecules of hydrogen chloride (HCl) than between molecules of hydrogen fluoride (HF).




3. The sulfur atom is larger than the chlorine atom (ie it has a greater covalent radius – 0.104 nm compared to 0.099 nm).



4. Chlorine is less electronegative than fluorine.




5. Arsine ( $\text{AsH}_3$ ) is a weaker base than ammonia ( $\text{NH}_3$ ).







6. Potassium fluoride has a smaller lattice energy ( $813 \text{ kJmol}^{-1}$ ) than sodium fluoride ( $915 \text{ kJmol}^{-1}$ ).



7. Less energy is released when sodium ions are hydrated ( $390 \text{ kJmol}^{-1}$ ) than when lithium ions are hydrated ( $499 \text{ kJmol}^{-1}$ ).



8. Chlorine is more electronegative than sulfur.



# Explaining chemical phenomena (1)

Chemists use their models and theories to try and explain phenomena about chemical systems. Suggest an explanation for each of the following (you may find it useful to refer to a periodic table):

1. Lithium has a higher melting temperature (454 K) than sodium (371 K).

---

---

---

---

---

2. There is stronger bonding, called hydrogen bonding, between molecules of water ( $\text{H}_2\text{O}$ ) than between molecules of hydrogen sulfide ( $\text{H}_2\text{S}$ ).

---

---

---

---

---

3. The nitrogen atom is smaller than the carbon atom (*ie* it has a smaller covalent radius – 0.074 nm compared to 0.077 nm).

---

---

---

---

---

4. Chlorine is more electronegative than bromine.

---

---

---

---

---

5. Ammonia ( $\text{NH}_3$ ) is a stronger base than phosphine ( $\text{PH}_3$ ).

---

---

---

---

---

6. Magnesium chloride has a larger lattice energy ( $2489 \text{ kJmol}^{-1}$ ) than calcium chloride ( $2197 \text{ kJmol}^{-1}$ ).

---

---

---

---

---

7. More energy is released when sodium ions are hydrated ( $390 \text{ kJmol}^{-1}$ ) than when potassium ions are hydrated ( $305 \text{ kJmol}^{-1}$ ).

---

---

---

---

---

8. Nitrogen is less electronegative than oxygen.

---

---

---

---

---

# Constructing chemical explanations

Chemists use models and theories to try and explain phenomena about chemical systems. Although chemists seems to use a wide range of different models and theories, many of them are based on the same few basic principles.

If you can learn about these basic ideas you can use them as 'tools' to build up chemical explanations.

## Some basic ideas used in explaining chemistry

### The importance of size and charge

A large number of chemical phenomena can be explained, at least partly, in terms of simple ideas like the size and charge on ions or other particles.

### Charge density

If two ions have the same charge, but are different in size, then the smaller one is said to have a greater density of charge. The ion with the greater density of charge can often have a greater effect – if it can get up close to other ions or molecules.

An ion with greater charge density can form a stronger bond with an oppositely charged ion. This can lead to a more stable ionic lattice, which therefore requires more energy to disrupt. (So the lattice energy of magnesium chloride is greater than the lattice energy of calcium chloride - as the  $Mg^{2+}$  ion, which is smaller, has a greater charge density than the  $Ca^{2+}$  ion, even though they have the same charge.)

If the ions are of similar size, but have a different charge the one with the greater charge will have the larger charge density, and may be able to form stronger bonds.

When ionic materials dissolve in water the ions are hydrated (surrounded by water molecules which bond to them). The greater the charge density of an ion the more water molecules will bond to it, and the more energy will be released when a material with that type of ion dissolves.

These ideas do not always help us predict what will happen in experiments, as sometimes there are several effects operating at once. For example, an ion with a greater charge density can be hydrated more (which would make the material more soluble) but will usually bond more strongly to oppositely charged ions (which would make the material harder to dissolve!)

In some books the terms charge:size (charge to size ratio) or charge:volume (charge to volume ratio) may be used instead of charge density.

### Core charge

Many chemical processes can be – at least partly – explained in terms of the charges in the ions or molecules involved. Atoms are neutral, but separate atoms are seldom involved in chemical processes. Usually we are concerned with ions or molecules.

In what ways are ions and molecules like atoms, and how are they different?

Atoms may be thought of as a positive nucleus surrounded by several shells of electrons. (Of course, the electronic structure is more complicated than that, with different types of orbitals. However, it is often useful to think in terms of shells.) Most of the time the nuclei of the atoms do not change (and when they do this is studied by physicists). Usually only the outermost shell of electrons, the valence shell, is changed in chemical processes. The nucleus and all the inner shells are usually not significantly changed.

The term core is used to describe the nucleus of an atom, plus all the electrons that are not in the outer (valence) shell.

The charge on an atomic core is called the core charge.

It is often useful to know what the core charge is. The core charge will equal the positive charge on the nucleus plus the negative charge of all the inner-shell electrons.



The table shows how to calculate core charge.

Complete the table.

Element E	Atomic Number Z	Number of protons $n_p$	Number of electrons $n_e$	Electronic configuration e.c.	Number of outershell electrons $n_{ve}$	Number of core electrons $n_{ce} (=n_e - n_{ve})$	To calculate core charge $n_p - n_{ce} =$	Core charge c.c.
H	1	1	1	1	1	0	$1 - 0 =$	+1
He	2	2	2	2	2	0	$2 - 0 =$	+2
Li	3	3	3	2.1	1	2	$3 - 2 =$	+1
Be	4	4	4	2.2	2	2	$4 - 2 =$	+2
B	5	5	5	2.3	3	2	$5 - 2 =$	+3
C	6	6	6	2.4		2		+4
N	7	7	7		5	2	$7 - 2 =$	+5
O	8	8	8	2.6	6	2	$8 - 2 =$	
F	9	9	9	2.7				+7
Ne	10	10	10	2.8			$10 - 2 =$	
Na	11	11	11	2.8.1	1	10	$11 - 10 =$	+1
Mg				2.8.2	2	10	$12 - 10 =$	+2
Al	13	13	13	2.8.3		10		+3
Si	14			2.8.4	4			+4
P				2.8.5		10		
S					6	10		
Cl				2.8.7				+7
Ar	18	18	18	2.8.8	8	10	$18 - 10 =$	+8
K	19			2.8.8.1	1	18	$19 - 18 =$	
Ca	20	20	20		2			+2

The diagram below shows the symbols for the first 20 elements. Using the table on the previous page, complete the diagram by adding the core charges that are not shown.

Can you spot any pattern?

Core charge for the first 20 elements							
H +1							He +2
Li +1	Be +2	B	C +4	N +5	O	F +7	Ne +8
Na +1	Mg +2	Al +3	Si +4	P	S	Cl +7	Ar +8
K	Ca +2						

## Electronegativity

One of the most useful chemical concepts is that of electronegativity.

Electronegativity is the tendency of an element to attract the bonding electrons towards itself in compounds.

The electronegativity of an element depends upon how strongly the outer (valence) shell electrons are attracted to the core. The greater the charge on the core, and the nearer the outer shell of electrons are, the more strongly they are bound to the core.

Electronegativity is therefore greater at the top of a group (where the outer shell of negatively charged electrons is pulled closest to the positively charged nucleus) and to the right of a period (where the core charge is greatest):


Approximate core charge density for some of the elements							
H ⊕ <sup>+1</sup>							
Li ⊕ <sup>+1</sup>	Be ⊕ <sup>+2</sup>	B ⊕ <sup>+3</sup>	C ⊕ <sup>+4</sup>	N ⊕ <sup>+5</sup>	O ⊕ <sup>+6</sup>	F ⊕ <sup>+7</sup>	
Na ⊕ <sup>+1</sup>	Mg ⊕ <sup>+2</sup>	Al ⊕ <sup>+3</sup>	Si ⊕ <sup>+4</sup>	P ⊕ <sup>+5</sup>	S ⊕ <sup>+6</sup>	Cl ⊕ <sup>+7</sup>	
K ⊕ <sup>+1</sup>	Ca ⊕ <sup>+2</sup>						





## Bond polarity in terms of electronegativity

Where the electrons in a bond are pulled equally by both cores, the bond is non-polar.



For example in  $F_2$  both the atoms of fluorine have cores of the same size and charge.



However, in the molecule of  $CF_4$  the fluorine atoms have a more highly charged core, and the bonding electrons are pulled closer to the fluorine cores than carbon core. The bonds are polar. This can be represented as:



The inter-halogen compound  $ClF$  also has a polar bond. Although the cores of both atoms have the same charge, the fluorine core has a higher charge density. This can be represented as:



## Building up chemistry explanations

Many of the explanations we construct in chemistry use ideas such as charge density, core charge and electronegativity. Many explanations can be built up using just a few basis ideas. Certain key phrases can be used as tools for building up explanations. (This does not mean you can just pick any key phrase: you have to understand the chemistry, and select the right phrases for a particular explanation!)

Consider the question: why do water molecules bond together?

Water molecules bond together because they are polar.

Water molecules are polar because the hydrogen-oxygen bond is polar.

The hydrogen-oxygen bond is polar because oxygen is more electronegative than hydrogen.

Oxygen is more electronegative than hydrogen because oxygen has a larger core charge.

This is not a 'complete' explanation. It would be possible to explain in more detail about the core charge (eg oxygen has a core charge of +6 because it has a nucleus of charge +8, partly shielded by 2 electrons in the core). It is also important to know that a molecule with polar bonds is not always a polar molecule! (So the molecule of  $\text{CF}_4$  discussed above has four polar bonds, but overall the molecule is non-polar.) The explanation above could be improved by adding:

... Water molecules are polar because the hydrogen-oxygen bond is polar, and the water molecule has two polar bonds which do not cancel out ...

This theme could also be continued...

... The two polar bonds do not cancel because the water molecule is angular.


The water molecule is angular because there are four electron pairs in the oxygen outer shell.

There are two non-bonding (lone) pairs as well as the two pairs of bonding electrons ...

So even a simple question could have a very complicated answer if we want to give a full and detailed explanation. You do not normally need to do this, although it can be good practice in testing how well you understand your chemistry.

## Charting your explanations


If some explanations seem rather complicated, you may find it helps to break the explanation down into steps, which can be put into a flow chart. In the example below, the phenomenon to be explained is written at the top of the box, and each arrow may be read as a 'because'.



Although such a scheme may seem complicated, it helps to learn to think about the individual steps in the explanations. If you can understand the steps you can learn to put together such schemes.

For example consider the question: why does water have a higher boiling temperature than would be predicted from its molecular mass?

You may know that it is because the hydrogen bonding in water holds the molecules together, so that more energy is needed to separate the molecules. In other words, we can amend the scheme above.



### Some key phrases

The following phrases are examples of some of those that may be useful in constructing explanations (you may want to add to this list):


- has strong(er)/weak(er) bonding
- is more/less electronegative
- has a higher/lower charge density
- has a larger/smaller core charge
- electrons are nearer/further from the nucleus

# Completing explanations

## 1. Explaining that lithium has a higher melting temperature than sodium

Complete the missing words in the scheme (left).

In your own words try and explain why lithium has a higher melting temperature (454 K) than sodium (371 K).




---



---



---



---



---



---



---



---



---



---



---



---



---



---



---




---

Can you redraw the scheme to include extra detail, and so make it more informative?





# Completing explanations



## 4. Explaining that chlorine is more electronegative than bromine

Complete the missing words in the scheme (left).

In your own words try and explain why chlorine is more electronegative than bromine.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

Can you redraw the scheme to include extra detail, and so make it more informative?



# Completing explanations



5. Explaining that ammonia (NH<sub>3</sub>) is a stronger base than phosphine (PH<sub>3</sub>)

Complete the missing words in the scheme (left).

In your own words try and explain why ammonia (NH<sub>3</sub>) is a stronger base than phosphine (PH<sub>3</sub>).

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Can you redraw the scheme to include extra detail, and so make it more informative?

# Completing explanations



Magnesium chloride has a larger lattice energy than calcium chloride

More energy is required to disrupt the \_\_\_\_\_ chloride lattice

The bonding is stronger in magnesium chloride than in calcium chloride

Chloride anions get closer to the centres of positive charge in \_\_\_\_\_ chloride

Magnesium 2+ cations are \_\_\_\_\_ than calcium 2+ cations

Magnesium is in period 3 and calcium is in period 4

6. Explaining that magnesium chloride has a larger lattice energy ( $2489 \text{ kJmol}^{-1}$ ) than calcium chloride ( $2197 \text{ kJmol}^{-1}$ )

Complete the missing words in the scheme (left).

In your own words try and explain why magnesium chloride has a larger lattice energy than calcium chloride.

---

---

---

---

---

---

---

---

---


---

---

Can you redraw the scheme to include extra detail, and so make it more informative?



# Completing explanations



7. Explaining that more energy is released when sodium ions are hydrated ( $390 \text{ kJmol}^{-1}$ ) than when potassium ions are hydrated ( $305 \text{ kJmol}^{-1}$ )

Complete the missing words in the scheme (left).

In your own words try and explain why more energy is released when sodium ions are hydrated than when potassium ions are hydrated.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Can you redraw the scheme to include extra detail, and so make it more informative?

# Completing explanations

8. Explaining that nitrogen is less electronegative than oxygen.

Complete the missing words in the scheme (left).

In your own words try and explain why nitrogen is less electronegative than oxygen.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

---

---

Can you redraw the scheme to include extra detail, and so make it more informative?



# Explaining chemical phenomena (2)

Chemists use their models and theories to try and explain phenomena about chemical systems. Suggest an explanation for each of the following (you may find it useful to refer to a Periodic Table).

1. Potassium has a lower melting temperature (336 K) than sodium (371 K).

---

---

---

---

---

2. There is weaker bonding between molecules of hydrogen chloride (HCl) than between molecules of hydrogen fluoride (HF).

---

---

---

---

---

3. The sulfur atom is larger than the chlorine atom (*ie* it has a greater covalent radius – 0.104 pm compared to 0.099 pm).

---

---

---

---

---

4. Chlorine is less electronegative than fluorine.

---

---

---

---

---

5. Arsine ( $\text{AsH}_3$ ) is a weaker base than ammonia ( $\text{NH}_3$ ).

---

---

---

---

---

6. Potassium fluoride has a smaller lattice energy ( $813 \text{ kJmol}^{-1}$ ) than sodium fluoride ( $915 \text{ kJmol}^{-1}$ ).

---

---

---

---

---

7. Less energy is released when sodium ions are hydrated ( $390 \text{ kJmol}^{-1}$ ) than when lithium ions are hydrated ( $499 \text{ kJmol}^{-1}$ ).

---

---

---

---

---

8. Chlorine is more electronegative than sulfur.

---

---

---

---

---

